


Part I - Ageing

Sex, Ageing and Foraging Theory

What is ageing?

aka senescence


- Gradual deterioration of function.




What is ageing?

aka senescence


- Gradual deterioration of function.
- Decrease in survival rate and/or fecundity with age.




Trends in Cell Biology 2020 30777-791DOI: (10.1016/j.tcb.2020.07.002)




Natural variation in ageing and lifespan



Natural variation in ageing and lifespan




Treaster S, Karasik D and Harris MP (2021 Front. Genet. 12:678073.doi: 10.3389/fgene.2021.678073



Homo sapiens



Gopherus agassizii
(desert tortoise)




Jones *et al. Nature* 000, 1-5 (2013) doi:10.1038/nature12789

Modelling age structure

Dynamics of an age-structured population

The Demography of the World Population from 1950 to 2100

Shown is the age distribution of the world population – by sex – from 1950 to 2018 and the *UN Population Division's* projection until 2100.




Data source: United Nations Population Division – World Population Prospects 2017; Medium Variant.
The data visualization is available at [OurWorldinData.org](https://ourworldindata.org), where you find more research on how the world is changing and why.

Licensed under CC-BY by the author Max Roser.


Dynamics of an age-structured population

- $n_{a,t}$ = n. of individuals of age a at time t
- p_a = probability of survival from age a to $a+1$
- m_a = fecundity at age a (i.e. number of newborns)
- $f_a = p_0 m_a$ = effective fecundity at age a (i.e. number newborns that survive to age 1, with probability p_0)



Dynamics of an age-structured population


$$n_{1,t+1} =$$



Dynamics of an age-structured population

$$n_{1,t+1} = \sum_{a=1}^A f_a n_{a,t}$$

\swarrow
 $p_0 m_a$




Dynamics of an age-structured population

$$n_{1,t+1} = \sum_{a=1}^A f_a n_{a,t}$$

\swarrow $p_0 m_a$

$$n_{a+1,t+1} =$$




Dynamics of an age-structured population

$$n_{1,t+1} = \sum_{a=1}^A f_a n_{a,t}$$

\swarrow
 $p_0 m_a$

$$n_{a+1,t+1} = p_a n_{a,t} \text{ for } a = 1, 2, \dots, A-1$$



Leslie Matrix

$$(A\mathbf{v})_j = \sum_i a_{ij}v_i$$

$$(AB)_{ik} = \sum_j a_{ij}b_{jk}$$

$$n_{1,t+1} = \sum_{a=1}^A f_a n_{a,t}$$

\swarrow
 $p_0 m_a$

$$n_{a+1,t+1} = p_a n_{a,t} \text{ for } a = 1, 2, \dots, A-1$$

$$\mathbf{n}_t = \begin{pmatrix} n_{1,t} \\ n_{2,t} \\ n_{3,t} \\ \vdots \\ n_{A-1,t} \\ n_{A,t} \end{pmatrix}$$

$$\mathbf{L} = \begin{pmatrix} f_1 & f_2 & f_3 & \dots & f_{A-1} & f_A \\ p_1 & 0 & 0 & \dots & 0 & 0 \\ 0 & p_2 & 0 & \dots & 0 & 0 \\ 0 & 0 & p_3 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & p_{A-1} & 0 \end{pmatrix}$$

$$\mathbf{n}_{t+1} = \mathbf{L} \mathbf{n}_t$$

Asymptotic behaviour

$$n_{t+1} = L n_t$$

$$n_1 = L n_0$$

$$n_2 = L n_1 = L^2 n_0$$

$$n_3 = L n_2 = L^3 n_0$$

$$\vdots$$

$$n_t = L^t n_0$$

Asymptotic behaviour

$$n_{t+1} = L n_t$$

$$n_1 = L n_0$$

$$n_2 = L n_1 = L^2 n_0$$

$$n_3 = L n_2 = L^3 n_0$$

⋮

$$n_t = L^t n_0$$

Age a (years)	p_a	m_a	f_a
0	0.25		
1	0.46	1.28	0.32
2	0.77	2.28	0.57
3	0.65	2.28	0.57
4	0.67	2.28	0.57
5	0.64	2.28	0.57
6	0.88	2.28	0.57
7		2.28	0.57

$$L = \begin{pmatrix} 0.32 & 0.57 & 0.57 & 0.57 & 0.57 & 0.57 & 0.57 \\ 0.46 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0.77 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0.65 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0.67 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0.64 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0.88 & 0 \end{pmatrix}$$



Exponential increase

$$n_{t+1} = L n_t$$

$$n_1 = L n_0$$

$$n_2 = L n_1 = L^2 n_0$$


$$n_3 = L n_2 = L^3 n_0$$

$$\vdots$$

$$n_t = L^t n_0$$

Age <i>a</i> (years)	<i>p_a</i>	<i>m_a</i>	<i>f_a</i>
0	0.25		
1	0.46	1.28	0.32
2	0.77	2.28	0.57
3	0.65	2.28	0.57
4	0.67	2.28	0.57
5	0.64	2.28	0.57
6	0.88	2.28	0.57
7		2.28	0.57

$$L = \begin{pmatrix} 0.32 & 0.57 & 0.57 & 0.57 & 0.57 & 0.57 & 0.57 \\ 0.46 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0.77 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0.65 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0.67 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0.64 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0.88 & 0 \end{pmatrix}$$



Extinction

$$n_{t+1} = L n_t$$

$$n_1 = L n_0$$


$$n_2 = L n_1 = L^2 n_0$$

$$n_3 = L n_2 = L^3 n_0$$

⋮

$$n_t = L^t n_0$$

Age a (years)	p_a	m_a	f_a
0	0.25 0.2		
1	0.46	1.28	0.256
2	0.77	2.28	0.456
3	0.65	2.28	0.456
4	0.67	2.28	0.456
5	0.64	2.28	0.456
6	0.88	2.28	0.456
7		2.28	0.456



Stable age distribution


$$n_{t+1} = L n_t$$

$$\begin{aligned} n_1 &= L n_0 \\ n_2 &= L n_1 = L^2 n_0 \\ n_3 &= L n_2 = L^3 n_0 \\ &\vdots \\ n_t &= L^t n_0 \end{aligned}$$

Age a (years)	p_a	m_a	f_a
0	0.25		
1	0.46	1.28	0.32
2	0.77	2.28	0.57
3	0.65	2.28	0.57
4	0.67	2.28	0.57
5	0.64	2.28	0.57
6	0.88	2.28	0.57
7		2.28	0.57

$$c_{a,t} = \frac{n_{a,t}}{\sum_{a=1}^A n_{a,t}}$$

= proportion of individuals of age a at time t




Growth rate

In the long run (large t),

$$\mathbf{n}_t \rightarrow c_0 \lambda^t \mathbf{u}$$

where:

- λ is the growth rate (and leading eigenvalue of the Leslie matrix L);
- \mathbf{u} is the stable age distribution (associated right eigenvector, i.e. $L\mathbf{u} = \lambda\mathbf{u}$, with entries summing to one);
- $c_0 = \mathbf{v} \cdot \mathbf{n}_0 > 0$ is a positive constant, where \mathbf{v} is vector of reproductive values (given by $\mathbf{v}^T L = \lambda \mathbf{v}$, such that $\mathbf{v}^T \mathbf{u} = 1$).




Reproductive values

In the long run (large t),

$$\mathbf{n}_t \rightarrow c_0 \lambda^t \mathbf{u}$$

where:

- λ is the growth rate (and leading eigenvalue of the Leslie matrix L);
- \mathbf{u} is the stable age distribution (associated right eigenvector, i.e. $L\mathbf{u} = \lambda\mathbf{u}$, with entries summing to one);
- $c_0 = \mathbf{v} \cdot \mathbf{n}_0 > 0$ is a positive constant, where \mathbf{v} is vector of reproductive values (given by $\mathbf{v}^T L = \lambda \mathbf{v}$, such that $\mathbf{v}^T \mathbf{u} = 1$).



reproductive value ~ relative importance of individuals of different ages in the initial population in determining the total population size in the distant future



Explosion vs. Extinction

In the long run (large t),

$$\mathbf{n}_t \rightarrow c_0 \lambda^t \mathbf{u}$$

where:

- λ is the growth rate (and leading eigenvalue of the Leslie matrix L);
- \mathbf{u} is the stable age distribution (associated right eigenvector, i.e. $L\mathbf{u} = \lambda\mathbf{u}$, with entries summing to one);
- $c_0 = \mathbf{v} \cdot \mathbf{n}_0 > 0$ is a positive constant, where \mathbf{v} is vector of reproductive values (given by $\mathbf{v}^T L = \lambda \mathbf{v}$, such that $\mathbf{v}^T \mathbf{u} = 1$).



Population grows exponentially at rate λ when $\lambda > 1$ (otherwise goes extinct when $\lambda < 1$).

Age distribution stabilises to \mathbf{u} .



Lifetime reproductive success

$$R_0 = \sum_{a=1}^A l_a m_a$$

$l_a = p_0 p_1 p_2 \dots p_{a-1}$ = probability of survival until age a

= lifetime reproductive success

= expected number of offspring during one's lifetime.




$\lambda > 1$ if and only if $R > 1$

Density-dependence

- Competition for resources —> density regulation
- i.e. survival and/or reproduction decreases with population size
- Leslie matrix now depends on n_t $L(n_t)$
- Population size converges to equilibrium where $R_0 = 1$



Effective fecundity




Convergence to demographic equilibrium

- Competition for resources \rightarrow density regulation
- i.e. survival and/or reproduction decreases with population size
- Leslie matrix now depends on \mathbf{n}_t $L(\mathbf{n}_t)$
- Population size converges to equilibrium where $R_0 = 1$

$0.25 / \left(1 + \gamma \sum_{a=1}^A n_{a,t} \right)$


Age a (years)	p_a	m_a	f_a
0	0.25		
1	0.46	1.28	0.32
2	0.77	2.28	0.57
3	0.65	2.28	0.57
4	0.67	2.28	0.57
5	0.64	2.28	0.57
6	0.88	2.28	0.57
7		2.28	0.57



Evolution in age-structured population


Mutant fitness and reproductive success

- Consider a population monomorphic for trait x (e.g. bill size, neck length) at demographic equilibrium (under density-dependent regulation).



Mutant fitness and reproductive success


- Consider a population monomorphic for trait x (e.g. bill size, neck length) at demographic equilibrium (under density-dependent regulation).
- Suppose a mutant appears with alternative trait y .



Mutant fitness and reproductive success

- Consider a population monomorphic for trait x (e.g. bill size, neck length) at demographic equilibrium (under density-dependent regulation).
- Suppose a mutant appears with alternative trait y .

Is the mutant going to invade and replace the resident ?




Mutant fitness and reproductive success

- Consider a population monomorphic for trait x (e.g. bill size, neck length) at demographic equilibrium (under density-dependent regulation).
- Suppose a mutant appears with alternative trait y .
- In a large well mixed population, mutant invades only if

$$R_0(y, x) = \sum_{a=1}^A l_a(y, x) m_a(y, x) > 1$$

Pr of survival to
age a of a rare y
mutant in a
population of x

Fecundity of a
rare y mutant of
age a in a
population of x




Mutant fitness and reproductive success

- Consider a population monomorphic for trait x (e.g. bill size, neck length) at demographic equilibrium (under density-dependent regulation).
- Suppose a mutant appears with alternative trait y .
- In a large well mixed population, mutant invades only if


$$R_0(y, x) = \sum_{a=1}^A l_a(y, x) m_a(y, x) > 1$$

i.e. if a mutant on average has more than one offspring over its lifetime.




Evolutionary analysis

- For quantitative traits, mutant lifetime reproductive success $R_0(y, x)$ defines a **fitness landscape**.



Evolutionary analysis



- For quantitative traits, mutant lifetime reproductive success $R_0(y, x)$ defines a **fitness landscape**.
- An evolving population **climbs** this landscape to arrive to a maximum.



Evolutionary analysis

- For quantitative traits, mutant lifetime reproductive success $R_0(y, x)$ defines a **fitness landscape**.
- An evolving population **climbs** this landscape to arrive to a maximum.
- The direction of this climb is given by the **selection gradient**,

$$s(x) = \left. \frac{\partial R(y, x)}{\partial y} \right|_{y=x}$$



Evolutionary analysis

- For quantitative traits, mutant lifetime reproductive success $R_0(y, x)$ defines a **fitness landscape**.
- An evolving population **climbs** this landscape to arrive to a maximum.
- The direction of this climb is given by the **selection gradient**,



$$s(x) = \left. \frac{\partial R(y, x)}{\partial y} \right|_{y=x}$$

- A maximum x^* is such that

$$s(x^*) = 0$$


and

$$\left. \frac{\partial s(x)}{\partial x} \right|_{x=x^*} < 0$$



Summary

- Dynamics of age structured population modelled via the Leslie matrix.
- The population grows when lifetime reproductive success R_0 is above 1.
- Due to competition, natural populations eventually experience density-dependent competition.
- Populations thus stabilise to a demographic equilibrium where $R_0 = 1$.
- A rare mutant y invades an x population at demographic equilibrium when mutant reproductive success $R_0(y, x) > 1$.



$$R_0 = \sum_{a=1}^A l_a m_a$$

